


Enjoy beautiful results for up to 18 months when you follow the *Restylane*® Regimen.

Bring this form to your healthcare professional to create your personalized *Restylane*® Regimen. At your initial treatment, you and your healthcare professional may want to set up follow-up appointments to help you achieve correction for up to 18 months.

This is a suggested course of treatment. Consult your healthcare professional to develop your individual treatment options.



1 Ask your healthcare professional how much *Restylane*® he or she recommends for your initial treatment to achieve full correction.

2 Make an appointment for a follow-up treatment at 4.5 or 9 months later.

3 Work with your healthcare professional to maintain a *Restylane*® Regimen that's right for you.

Notes _____

Notes _____

Notes _____

See Important Safety Considerations on following page.





Important Safety Considerations of *Restylane*[®]

Restylane[®] restores volume and fullness to the skin to correct moderate to severe facial wrinkles and folds, such as the lines from your nose to the corners of your mouth (nasolabial folds). After your treatment, you might have some swelling, redness, pain, bruising, and tenderness. This will normally last less than seven days. Although rare, red or swollen small bumps may occur. If you have had facial cold sores before, an injection can cause another outbreak. In rare circumstances, the doctor may inject into a blood vessel, which can damage the skin. To avoid bruising and bleeding, you should not use *Restylane*[®] if you have recently used drugs that thin your blood or prevent clotting. If you are pregnant, breastfeeding, or under 18 years, you should not use *Restylane*[®].

Restylane[®] should not be used by people with previous bad allergies, particularly to certain microorganisms known as gram-positive bacteria; by people with previous bad allergies to drugs that have required in-hospital treatment; or by people with bleeding disorders. *Restylane*[®] should not be injected anywhere except the skin or just under the skin.

The use of *Restylane*[®] at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. Use of *Restylane*[®] in these instances could delay healing or make your skin problems worse.

Restylane[®] is available only through a licensed practitioner. Complete product and safety information is available at www.RestylaneUSA.com.

